

THE ANTIFUNGAL COATING OF DIFFERENT WALL FINISHING
TOWARDS GROWTH INHIBITION OF *ASPERGILLUS NIGER*

UMI KALTHSOM BINTI PARJO

A thesis submitted in
fulfillment of the requirements for the award of the
Doctor of Philosophy in Engineering Technology



Faculty of Engineering Technology
Universiti Tun Hussein Onn Malaysia

JULY 2018

DEDICATION

Specially dedicated to my beloved mother, Hajah Sa'diah binti Misman
and to my late father Haji Parjo bin Singo, brothers and sisters,
my supervisor Assoc. Prof. Dr. Norshuhaila binti Mohamed Sunar
and co-supervisor Assoc. Prof. Dr. Abdul Mutalib bin Leman
my husband Akmal Hakim bin Remle
and to all my friends especially, Nurulliza binti Mohamad, Yusma Yusnida binti
Yusak, Fatin Malina binti Mukamat Alip and Noor Izzah binti Kamin
for your endless support.



PTTA UTHM
PERPUSTAKAAN TUNKU TUN AMINAH

ACKNOWLEDGEMENTS

Alhamdulillah, all praise to Allah, the Most Merciful and Most Wise for His SWT blessing and guidance, I could finish this thesis.

Firstly, I would like to express my deepest and sincere gratitude to my supervisor Assoc. Prof. Dr. Norshuhaila binti Mohamed Sunar for her valuable ideas, concern, help, dedication, suggestion, guidance, and her support either moral or physical to help me in completing this report. I'm also would like to forward my deepest and sincerely gratitude towards my co-supervisor Assoc. Prof. Dr. Abdul Mutalib bin Leman for all his comments, guidance, support and sharing knowledge throughout this research. Without them, I will be lost in pursuing my PhD.

My special appreciation goes to the science officers and technicians of microbiology and environmental analysis laboratory for their kindness and assistance during the experimental works. Special gratitude to all my friends whose get involved in this project for their cooperation, help, and support during the laboratory works.

Lastly, to my beloved mother and father, brothers and sisters who have supported me to complete this project. Without their prayers, encouragement, and understanding it would be difficult to me to finish this work. Special thanks to MYPhD 15, for sponsoring me throughout 3 years my study at UTHM. Finally, to my entire friend who have brought the best memories throughout the whole process in doing this study, thanks you for everything and may Allah bless all of you.

ABSTRACT

Indoor air quality (IAQ) is an essential factor which directly affects human and the comfort of building occupants. The concentration of fungi, humans, animals and others contribute to indoor air pollutants and can cause a syndrome known as sick building syndrome. In this study, the growth of fungi in a lecturers' room at Universiti Tun Hussein Onn Malaysia (UTHM) was investigated with the application of three types of antifungals namely potassium sorbate, calcium benzoate and zinc salicylate. The study was conducted to enumerate and evaluate the growth of *Aspergillus niger* (*A. niger*). Four different types of wall finishings were utilized; namely acrylic paint, glycerol-based paint, thin wallpaper and thick wallpaper. These wall finishings were applied after being incorporated with antifungals on three different building materials; namely plasterboard, concrete and wood. Each sample was placed with 50 μ l of the spore suspension on a petri-dish containing malt extract agar. The sample was incubated at the relative humidity and temperature of 90% and 32°C respectively, based on the results of preliminary work. A visual assessment was made according to ASTM D5590-00 (2012). The results of antifungal coating resistance test showed that the best antifungal to remediate *A. niger* is potassium sorbate applied on thick wallpaper surfaces for plasterboard with growth percentages of about 47%. However, in the assessment of the inhibitory activity of *A. niger*, the results showed that concrete treated by zinc salicylate performed better than other antifungals used to inhibit *A. niger* with 50% on thick wallpaper covering. From this investigation, it was shown that the potassium sorbate antifungal was effective in decreasing the growth rate of the *A. niger*. However, the effectiveness of the results also depended on the construction building materials used. This study provides a reference in the inhibition of *A. niger* using different antifungals in various types of building materials. In addition, the effectiveness of antifungal compounds used in this study also exhibited enormous potential to be used as additives to prevent indoor fungal growth in Malaysia's construction industry.

ABSTRAK

Kualiti udara dalaman (IAQ) adalah penting untuk kesihatan dan keselesaan penghuni bangunan. Kepekatan kulat daripada udara, manusia, binatang dan lain-lain menjadi punca kepada sindrom yang dikenali sebagai bangunan sakit dan pencemaran udara dalaman. Dalam kajian ini, pertumbuhan kulat di bilik pensyarah, UTHM disiasat dan tiga jenis antikulat iaitu *potassium sorbate*, *calcium benzoate* dan *zinc salicylate* diaplikasikan. Kajian dijalankan untuk menghitung dan menilai pertumbuhan *Aspergillus niger* (*A. niger*) selepas menggabungkan antikulat pada empat kemasan dinding iaitu *acrylic paint*, *glycerol-based paint*, kertas dinding nipis dan kertas dinding tebal. Kemasan dinding yang telah digabungkan bersama antikulat diletakkan di atas permukaan bahan binaan yang berbeza iaitu *plasterboard*, konkrit dan kayu. 50ul *spore suspension* telah diletakkan di atas permukaan bahan binaan, diletakkan di dalam *petri-dish* yang mengandungi *malt extract agar*. Semua sample dieramkan pada kelembapan 90% suhu 32°C, di mana data kelembapan dan suhu berpandukan optimum faktor pada kajian awalan. Penilaian secara visual adalah berdasarkan standard ASTM D5590-00 (2012). Keputusan ujian *coating resistance*, antikulat terbaik untuk mengurangkan pertumbuhan *A. niger* adalah *potassium sorbate* pada permukaan *plasterboard* yang disaluti kertas dinding tebal sebanyak 47%. Walau bagaimanapun, bagi peratus aktiviti perencatan *A. niger* terbantut pada permukaan konkrit disaluti kertas dinding tebal dengan 50% selepas dirawat dengan antikulat *zinc salicylate*. Antikulat *potassium sorbate* dan *zinc salicylate* dapat mengurangkan kadar pertumbuhan *A. niger* bergantung kepada jenis bahan binaan yang digunakan di dalam pembinaan pembangunan. Kajian ini dapat menyediakan rujukan dalam perencatan *A. niger* dengan menggunakan antikulat melalui ujian *coating resistance*. Selain itu penggunaan antikulat yang berkesan di dalam kajian ini mempunyai potensi untuk digunakan sebagai bahan tambahan untuk mencegah pertumbuhan kulat dalaman di dalam industri pembinaan di Malaysia.

CONTENTS

TITLE	i
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENT	iv
ABSTRACT	v
ABSTRAK	vi
CONTENTS	vii
LIST OF TABLES	xi
LIST OF FIGURES	xiii
LIST OF SYMBOLS ABBREVIATION	xvi
LIST OF APPENDICES	xvii
CHAPTER 1	INTRODUCTION
	1
1.1	Introduction
	1
1.2	Background of the study
	1
1.3	Problem statement
	4
1.4	Research question
	6
1.5	Objective of study
	6
1.6	Scope of study
	7
1.7	Significance of study
	7
1.8	Thesis overview
	8
CHAPTER 2	LITERATURE REVIEW
	9
2.1	Introduction
	9
2.2	Indoor environmental quality (IEQ) and indoor air quality (IAQ)
	10
2.3	Sources of IAQ contaminant
	11
2.4	Parameters of IAQ status
	12

2.4.1	Temperature	12
2.4.2	Relative humidity	13
2.4.3	Air velocity	15
2.5	Classification of indoor air contaminations	15
2.5.1	Chemical contaminations	16
2.5.1.1	Carbon monoxide	16
2.5.1.2	Formaldehyde	17
2.5.1.3	Respirable particulates	18
2.5.1.4	Total volatile organic compounds	18
2.5.2	Biological contaminations	18
2.5.2.1	Fungi	19
2.5.2.2	Bacteria	23
2.6	Health effects due to poor IAQ	25
2.6.1	Health effects of ETS	26
2.6.2	Sick-building syndrome	27
2.6.3	Building related illness	29
2.7	Control of IAQ	29
2.7.1	Antifungals	30
2.7.1.1	Potassium sorbate	36
2.7.1.2	Calcium benzoate	38
2.7.1.3	Zinc salicylate	40
2.7.2	Construction material indoor fungal growth in building	41
2.7.2.1	Plasterboard	41
2.7.2.2	Woods	43
2.7.2.3	Concrete	45
2.7.3	Characteristic wall finishing	47
2.8	Standard guideline (IAQ) Malaysia	49
2.9	Summary	52
CHAPTER 3	RESEARCH METHODOLOGY	54
3.1	Introduction	54
3.2	Experimental and method development	54
3.3	Indoor air quality (IAQ) parameter	56

3.4	Indoor fungal sampling	57
3.4.1	Enumeration of sample	60
3.4.2	Indoor fungal isolation	60
3.4.3	Indoor fungal identification	61
3.5	Indoor fungal suspension spore	61
3.6	Growth optimization of isolated <i>A. niger</i>	63
3.6.1	Temperature	63
3.6.2	Relative humidity	64
3.7	Optimization of growth measurement by Response Surface Methodology (RSM)	65
3.8	Building construction material coating resistance test	66
3.8.1	Characteristic of paint	69
3.8.2	<i>A. niger</i> growth percentage observation technique	69
3.9	Inhibition performance of <i>A. niger</i> growth	71
3.10	Safety procedures	73
3.11	Summary	74
CHAPTER 4	RESULT AND DISCUSSION	76
4.1	Introduction	76
4.2	Indoor air quality inspection and data collection	76
4.3	Enumeration of indoor sample	79
4.4	Optimization of temperature and relative humidity of the <i>A. niger</i> growth	81
4.5	Validation of Response Surface Methodology	83
4.6	Analysis of Coating Resistance Test	90
4.6.1	Coating resistance test on different substrates with various wall coverings	90
4.6.1.1	<i>A. niger</i> growth on plasterboard substrates	90
4.6.1.2	<i>A. niger</i> growth on concrete substrates	95
4.6.1.3	<i>A. niger</i> growth on wood substrates	100
4.6.2	Inhibitory activity of <i>A. niger</i> growth in application of antifungals	104

4.6.2.1 Inhibitory activity <i>A. niger</i> growth on plasterboard substrates	105
4.6.2.2 Inhibitory activity <i>A. niger</i> growth on concrete substrates	108
4.6.2.3 Inhibitory activity <i>A. niger</i> growth on wood substrates	111
4.7 Summary	116
CHAPTER 5 CONCLUSION AND RECOMMENDATIONS	117
5.1 Introduction	117
5.2 Conclusion	117
5.3 Recommendations and further works	119
REFERENCES	121
APPENDICES	142
VITA	



LIST OF TABLES

2.1	Relative humidity values for growth of variation fungal (Grant <i>et al.</i> , 1989; Brischke & Rapp, 2008)	15
2.2	Critical health effect from exposure to the formaldehyde (Nikulin <i>et al.</i> , 1994)	17
2.3	Studies on airborne fungi from the several countries	20
2.4	Disease syndromes associated with exposure to common fungal (IEH, 1996)	26
2.5	The health problems associated with moldy and damp building (Nielsen, 2002)	26
2.6	Most common symptoms of sick building syndrome (Jones, 1999)	28
2.7	Schematic review of the antifungal application purposes	33
2.8	The summaries of the potassium sorbate	36
2.9	Potassium sorbate for several antifungal purposes	37
2.10	Summary of the calcium benzoate application for antifungal purposes	39
2.11	Summary of the zinc salicylate application for antifungal purposes	40
2.12	Microorganisms identified on plasterboard surfaces in indoor environment	43
2.13	The indoor fungal in deterioration of wood material	44
2.14	The fungi in deterioration of concrete	46
2.15	The acceptable range for specific physical parameter according ICOP-IAQ by DOSH (2010)	50

2.16	The list of indoor air contaminants and the acceptable limits according to ICOP-IAQ by DOSH (2010)	50
3.1	Performance profile for bioaerosol sampling followed NIOSH Manual Analytical Method NAM0800 (1998)	59
3.2	The temperatures monitored at RH of 70% with respect to the incubation periods (days)	64
3.3	The relative humidities monitored at temperature at 26°C with respect to the incubation periods (days)	65
3.4	The levels of chosen variables	66
3.5	The degrees of indoor fungal growth (ASTM D5590-00)	69
3.6	Common ingredient component of paints	69
3.7	Chemical composition in emulsion of paints	70
3.8	Chemical composition in gloss paints	70
3.9	Summary of substrates used in study	72
3.10	Test, method, and equipment used to conduct the whole laboratory work	72
3.11	Summary of the safety precautions in the laboratory works	74
4.1	Summary result of IAQ parameters in room and comparison to DOSH (2010) Standard Guideline	79
4.2	Regression analysis (ANOVA) of quadratic polynomial model and significance test – Temperature	84
4.3	Regression analysis (ANOVA) of quadratic polynomial model and significance test – Relative humidity	85

LIST OF FIGURES

2.1	Potassium sorbate antifungal utilized in this study	31
2.2	Calcium benzoate antifungal utilized in this study	31
2.3	Zinc salicylate antifungal utilized in this study	32
2.4	The process of IAQ complaints and investigation according to DOSH, (2010)	52
3.1	The research flow of methodology	55
3.2	The bioaerosol pump (QuickTake® 30) equipment that were used for air sampling fungi collection.	59
3.3	The morphology of <i>A. niger</i> (40X) observed under light compound microscope	60
3.4	Method to prepare the spore suspension (a) Material used such as loop to transfer spore (b) loop was transferred in the flame (c) Then transfer into spore suspension (d) Duplicated spore suspension prepared (e) Calculation of spore suspension concentration by Microscope (Olympus CX300) (f) spore suspension was inoculated onto media	62
3.5	The number of visible <i>A.niger</i> cells calculated at the area of the Neubauer Chamber Improved	63
3.6	All the construction material substrates were soaked with 70% ethanol (a) plasterboard, (b) wood, (c) concrete	67
3.7	All the construction material substrates left to air dry before next process (a) plasterboard, (b) concrete, (c) woods	68
3.8	<i>A. niger</i> growth percentage observation and calculation technique according to ASTM D5590-00	71

4.1	The relative humidity measurement and its comparison with ICOP-IAQ by DOSH (2010)	77
4.2	Bar graph of temperature parameter and its comparison with DOSH ICOP-IAQ (2010)	78
4.3	The carbon dioxide measurement and its comparison with ICOP-IAQ by DOSH (2010)	79
4.4	The total indoor fungal count and its comparison with ICOP-IAQ by DOSH (2010)	80
4.5	The effect of temperature on the growth (mm) \pm SD of tested <i>A.niger</i>	82
4.6	The effect of relative humidity on the growth (mm) \pm SD of tested <i>A.niger</i>	83
4.7	Scatter diagram for experimental response versus predicted response temperature	86
4.8	Scatter diagram for experimental response versus predicted response relative humidity	86
4.9	Suggested in range value of factors for maximum response (growth temperature = 32°C)	87
4.10	Suggested in range value of factors for maximum response (growth relative humidity = 90%)	88
4.11	One factor plot showing the effect temperature on <i>A. niger</i> growth activity	88
4.12	One factor plot showing the effect relative humidity on <i>A.niger</i> growth activity	89
4.13	Bar graph showing the maximum desirability of 0.891 for the combined factors	89
4.14	The overall results for growth of <i>A. niger</i> growth on plasterboard substrate with different types of wall finishing as influenced by antifungals	92
4.15	The samples of substrates on plasterboard; (a) without application while (b) PS, (c) CB, (d) ZS are with application of antifungals	92

4.16	Summary of the <i>A. niger</i> growth on concrete with different types of wall finishing and different antifungals	97
4.17	The comparison between untreated and treated antifungals; (a) control samples (b)-(d) antifungals treated by PS, CB and ZS	97
4.18	The overall summary of <i>A. niger</i> for coating resistance test on a wood substrate with various surface coverings	102
4.19	The comparison substrates for control (a) and substrates treated with different antifungals (b) PS, (c) CB and (d) ZS	102
4.20	Overall graph inhibition of <i>A. niger</i> on treatment plasterboard substrates	105
4.21	Percentages of inhibition for each wall finishing treatment	105
4.22	(a) Spores actively appeared on the control samples surfaces (b) The effectiveness of PS to inhibit <i>A. niger</i> growth on thin wallpaper (THIN)	106
4.23	The inhibition of <i>A. niger</i> on treated concrete substrates	108
4.24	The antifungals inhibition on different wall finishing	108
4.25	The mycelium are seen on thick wallpaper (THICK) surfaces treated with zinc salicylate on (a) control samples (b) treatment sample	110
4.26	The inhibition of <i>A. niger</i> on treated wood substrates	111
4.27	The variation of antifungals in different wall finishing	111
4.28	SEM of hypha morphology (a) on wood control (b) covered by thick wallpaper (THICK) treated with the potassium sorbate	113
4.29	Schematic diagram of <i>A. niger</i> activity on the substrates	115
4.30	Schematic coating resistance test process	116

LIST OF SYMBOL AND ABBREVIATIONS

%	Percent
⁰ C	Degree Celsius
ASHRAE	American Society for Heating, Refrigerating and Air-Conditioning Engineers
ASTMD5590-00	American Society for Testing and Materials 5590-00
DOSH	Department of Occupational Safety and Health
EFSA	European Food Safety Authority
EPA	The United States Environmental Protection Agency
FKAAS	Faculty of Civil and Environmental Engineering
HVAC	Heating, Ventilation and Air Conditioning
ICOP-IAQ 2010	Industry Code of Practice on Indoor Air Quality 2010
IAQ	Indoor Air Quality
ISO 7730	International Standard Organization 7730
MEA	Malt Extracts Agar for Fungi
NIOSH	National Institute of Occupational Safety and Health
NMAM 0800	NIOSH Manual Analytical Method 0800
OSHA	Occupational Safety and Health Administration
RH	Relative Humidity
SBS	Sick Building Syndrome
TSA	Tryptocase Soy Agar
UTHM	Universiti Tun Hussein Onn Malaysia
VOC	Volatile Organic Compounds
WHO	World Health Organization

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Molecular biology result	142
B	Preparation sample for Coating Resistance Test	145
C	ANOVA results	148
D	Digital Microscopy Image Analyser (DMIA)	156
E	The observation results in antifungals treatment on different substrates	158
F	List of activities	179
G	Publications	182
H	Achievement	183

CHAPTER 1

INTRODUCTION

1.1 Introduction

Indoor air quality (IAQ) has become an important health topic since humans in general spend more than 80% of their time and activities in buildings. Humans may be exposed to various indoor air pollutants regardless of whether they are at their workplace or home (Wong *et al.*, 2008). IAQ which is caused by any indoor toxins, for example chemical contaminants (carbon monoxide, formaldehyde, ozone, respiratory particulates and total volatile organic compounds (TVOC), biological contaminants (bacterial and fungal), ventilation performance indicator (carbon dioxide) and any mass can cause significant health effects. Therefore, further investigation is required in order to prevent bio-deterioration on different building materials.

1.2 Background of the study

Contaminated of indoor air is common in all buildings and can cause the spread of infectious diseases by airborne spread. Exposure to fungi in the building can cause several types of human health problems, especially itching, infections, allergies and certain toxic effects. It has been shown that toxigenic organisms are the reason for the unfavorable impact on wellbeing (Shelton *et al.*, 2002) and contributes to a condition known as “sick building syndrome” (SBS). Therefore, it is imperative to discover an innovative approach to control or remediate the growth of indoor fungi.

There are five common contagious genera recognized in indoor air. They include *Cladosporium*, *Penicillium*, *Aspergillus*, *Alternaria*, and *Aureobasidium* (Vacher *et al.*, 2010; Bellotti *et al.*, 2013; Verdier *et al.*, 2014). The fungi species, *Cladosporium* and *Alternaria*, are mostly detected in outdoor air but have been detected in indoor air too. In addition, indoor contagious species such as, *Penicillium* and *Aspergillus* species can grow promptly on topsoil and rot litter (Hedayati, Mayahi, & Denning, 2010). Generally, fungi spread into the indoor environment through natural (open windows and doors) and mechanical ventilation systems (Gots, Layton, & Pirages, 2010; Laverge *et al.*, 2011). The indoor environment is influenced by a few variables including temperature, dampness, water seepage into structures, and the amount of open air entering the building (D'Orazio *et al.*, 2009; Deabes *et al.*, 2011; Alghamdi *et al.*, 2014).

In this study, the potential remediation of *Aspergillus niger* (*A. niger*) using a variation of antifungals has been explored. *A. niger* contains a toxin that can affect the health of those who work in the contaminated building. In addition, the effects of *A. niger* also contributes to sickness and allergic effects (Khan & Karuppayil, 2012; Flores *et al.*, 2013; Verdier *et al.*, 2014; Rakotonirainy, Vilmont, & Lavédrine, 2016).

The potential implications of indoor fungal growth to human health should not be underestimated; although the primary concern is the defacement of materials. The elimination and reduction of individual sources and their emissions are normally the effective method to control indoor air. Apart from that, the pollutants inside the building can be reduced when the amount of fresh air increases. In addition, humidity control helps to prevent moist air and decreases the growth of mould. However, humidity can be increased using a humidifier or decreased by opening the windows during dry weather (EPA, 2008).

The treatment of decay in buildings by synthetic antifungals has now been restricted due to their harmful effects on the environment, residual toxicity and carcinogenic nature (Khan & Karuppayil, 2012). Moreover, the constant use of chemicals has proven to induce resistance in a target organism. This concern has encouraged researchers to search for an efficient method to prevent and cure the harmful effects of microbes in a more eco-friendly manner (Sundis *et al.*, 2012). Thus, there is a need to find a method of controlling the growth of fungi approach to address the IAQ problems.

Many researchers have investigated indoor air quality, but only a few of them focused on remediation of fungal growth. Thus, further study on remediation of indoor fungal growth using different antifungals on building material is required. Bellotti *et al.* (2013) recognized the use of bioactive compounds from the food industry which have the capacity to control mould development in indoor waterborne coatings. In their studies, it was found that the addition of antifungals such as zinc salicylate and zinc benzoate proved that these antifungals inhibited the species of *Chaetomium globosum* and *Alternaria alternata*. In addition, Vacher *et al.* (2010) reported that the Dibromo-3-Nitrilopropionamide was used as antifungal to treat *Penicillium* or *Aspergillus*, *Chaetomium*, *Ulocladium*, *Stachybotrys* and *Cladosporium* on biodegradable materials. The antifungal nano zinc oxide and anatase titanium dioxide were found as the best photocatalytic agent and broadest spectrum of antimicrobial agents for interior paints to remediate *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Aspergillus niger* and *Penicillium chrysogenum* (Hochmannova & Vytrasova, 2010). Another study by Stanojevic *et al.* (2009) confirmed the potential of sodium benzoate, sodium nitrite, and potassium sorbate to reduce *Bacillus subtilis*, *Bacillus mycoides*, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Aspergillus flavus*, *Fusarium oxysporum*, *Candida albicans*, *Trichoderma harsianum* and *Penicillium italicum* and *Escherichia coli*.

Smilanick *et al.* (2008) highlighted the use of potassium sorbate, imazalil (imz), thiabendazole (tbz), pyrimethanil, and fludioxonil and found that the potassium sorbate was successful in remediating *Penicillium digitatum*. According to Clausen & Yang, (2007), potassium sorbate was found to inhibit the growth of about 5% for several types of fungus such as *Aspergillus niger*, *Penicillium chrysogenum*, and *Trichoderma viride* from outdoor sources. These studies showed that the antifungal compounds from various kinds of species have some potent antifungal ability.

However, the application of these antifungal in remediation of indoor microbial and fungal contamination has not been given full attention, especially in Malaysia. This is the motivation behind the present study. To best of the author's knowledge, no report was found on the evaluation of *A. niger* remediation using antifungals (potassium sorbate, calcium benzoate and zinc salicylate) applied on building materials. These antifungals were selected based on a previous study that showed that the antifungals which were applicable in controlling indoor fungi were mostly used in the food industry. However in this study, these antifungals is explored

explored for their use as antimicrobial additives for antimicrobial coatings in the construction industry, with concern placed on the safety of chemical used. The assessment of indoor air quality should be carried out and must take into consideration the safety of indoor occupants (DOSH, 2010). However, only several studies on indoor fungal remediation have been conducted at new buildings in this country. Rahman *et al.* (2012) conducted their investigation on the controlled growth of fungus in university's clinic in Malaysia but not in faculty buildings in Malaysia. Another study carried out in a secondary school building, found that fungal DNA and cat allergen were common in the Malaysian school (Halstensen, 2008).

Other researches to date tended to focus on others various species (Belloti *et al.*, 2013; Duangnapa, 2012; Deabes *et al.*, 2012) rather than that *A. niger* specifically. Therefore, the aim of the present study is to remediate the *A. niger* especially for common interior finishings of a building using several types of antifungals which include potassium sorbate, calcium benzoate and zinc salicylate that were previously used in food preservatives. It is hoped that this study would produce a new application to remediate the *A. niger* growth, after incorporating different wall finishings with antimicrobials coating. This was motivation for the present study.

1.3 Problem statement

Nowadays, issues concerning air quality with regard to indoor health environment are nothing new as they have been studied since 1850 during the hygienic revolution, and then followed by outdoor environmental issues (Spengler *et al.*, 2002). Similarly in Malaysia, indoor air quality has become prevalent recently to be explored. Although some buildings meet the recommended overall standard, there have been times that occupants would still complain about various parameters such as wall finishing and materials which are still not up to par with satisfying consumer needs. Thus, it is essential for the building to have a good indoor air quality (IAQ) as it affects the productivity and health of the occupants in the building. In fact, sustainable development results are not just in resource conservation only, but also in increasing productivity and occupant well-being (Syazwan *et al.*, 2009). More research needs to be conducted on issues regarding IAQ, especially on buildings in Malaysia with regard to the established standards that should be developed for the benefit to the occupants. Therefore, any IAQ issues related to residential, commercial, office and

institutional buildings where contamination has occurred by a variety of toxic or hazardous substance, as well as pollutants of biological origin requires serious solutions. This is due to the fact that microbial pollutants can cause the infection to the occupants in the building. For instance, fungus produces toxic mycotoxins, endotoxins, glucans and others in which are very harmful to human health (Verdier *et al.*, 2014).

Previous studies carried out in temperate region such as Japan's climate, with different periods of high temperatures and relative humidity were reasonable to the development of fungi (Nazar *et al.*, 2013). Similarly, Shelton *et al.* (2002) found that the occurrence of indoor fungus in the indoor environment was prevalent during the fall and summer seasons in the United States. According to Frankel *et al.* (2014), high relative humidity and temperature are among the factors that greatly contribute to the higher occurrence of indoor fungus. Additionally, others factor may influence indoor fungal growth, like the weather itself. For example, in Malaysia the weather is hot and humid during day and this leads to higher fungal growth at night. Furthermore, many of studies have shown that there is a relationship between indoor fungus and humidity to the health effect. This has also been shown to play a role in the production of mycotoxins (Vacher *et al.*, 2010). The temperature averages between 25°C and 35°C indicating a high indoor temperature, which is normally associated with local climate in Malaysia. However, this temperature range for the climate in Malaysia is in good agreement with DOSH (2010) where the best indoor temperature is supposed to be between 23°C and 26°C.

To date, there is still a lack in studies in Malaysia regarding the remediation of indoor fungus using antifungals in the construction industry. All the building materials currently used in construction, easily exposure to of mould growth especially in extreme moisture condition. The normal cleaning approaches however cannot provide an effective and sustainable solution to this mould problem. The continued replacement of damaged construction building materials with the new ones is neither cost-effective nor environmental friendly. This approach is not effective in Malaysia because of the high humidity throughout the year. Thus, there is a significantly need to search for sustainable, long term, low cost and environmental friendly approaches to address the IAQ problems. The issue of fungus in buildings is not new in Malaysia as it has been reported in the media and has occurred in hospitals, schools and various government buildings.

Recently, the UTHM authorities have received complaints of indoor fungal growth in the offices of academic buildings. As a result, UTHM's management has launched a research to find a lasting solution to this issue. The management is concerned that the fungus will affect the health of the lecturers and students. Therefore, the problem being addressed in this study is new as the application of the three selected antifungals (potassium sorbate, calcium benzoate and zinc salicylate) that are able to inhibit the growth of fungus in building materials (plasterboard, concrete and wood) are covered by different types of paint and wallpapers in Malaysian climate.

1.4 Research questions

This study was carried out to investigate the following research questions:

- 1) How the relative humidity and temperature levels influence the existence and growth of indoor fungal contamination?
- 2) What is the growth trend of *A. niger* when incorporated with optimum humidity of tropical climate?
- 3) How efficient the potassium sorbate, zinc salicylate and calcium benzoate antifungals used for coating bio-resistance test with different wall finishing can remediate or prevent *A. niger* on the substrates?

1.5 Objective of study

This study embarks on the following objectives:

- 1) To evaluate contamination of air in a room according to the Industry Code of Practice on Indoor Air Quality, 2010.
- 2) To optimize the specific growth rate of *A. niger* at different levels of temperature and relative humidity.
- 3) To develop a new application of coating resistance using antifungals in remediation specifically of *A. niger* on construction building materials covered with different wall finishing.
- 4) To compare the inhibition of *A. niger* growth after incorporation on three variations of treated wall finishing substrates.

1.6 Scope of study

- 1) This study used indoor fungal, *Aspergillus niger* that is isolated from a building. Selected antifungals consisted of potassium sorbate, calcium benzoate, and zinc salicylate, which were used to prevent indoor fungal growth from the air in the affected building.
- 2) The methods applied to collect indoor fungal are the single stage impactor (biosampler) in accordance with prerequisite of National Institute of Occupational Safety and Health (NIOSH) method (NIOSH Manual Analytical Method NMAM 0800).
- 3) Physical and chemical IAQ parameters were analyzed using guidelines of Industry Code of Practice of Indoor Air Quality in Malaysia (ICOP-IAQ) – DOSH, 2010.
- 4) The selected specific indoor fungus was inoculated at the concentration of 0.3×10^6 on substrates such as plasterboard, wood, and concrete.
- 5) In the coating resistance test, four wall finishing product were used on the plasterboard, concrete and wood surface which were acrylic paint, glycerol-based paint, thin wallpaper (80 g/m^2) and thick wall paper (240 g/m^2); and the indoor fungal growth was evaluated according to the ASTM D5590-00 (2012) standard scale.

1.7 Significance of study

The findings of this study contribute to the knowledge and understanding of the indoor fungal treatment using selected antifungals. This study is significant because:

- 1) The selections of the antifungal applied in this research have not been studied yet for indoor fungal treatment to remediate the *A. niger* that was isolated from indoor building especially in institutional buildings.
- 2) This study combines the function bio-coating resistance test using different wall finishing on different building material.
- 3) A new study of antifungal in combination with different wall finishing to remediate indoor fungal growth.
- 4) The selection of antifungal as antimicrobial additives for indoor fungal growth remediation can minimize the cost of renovation, demolition,

and reconstruction of sick building.

- 5) Provide an alternative prevention to avoid any hazardous to occupant and materials damage in building.

1.8 Thesis overview

This dissertation consists of five chapters. The chapter 1 describes the background of the study, problem statement, objectives, scope of study, and significance of study. Meanwhile chapter 2 explains a critical review of the research about indoor air quality, source of indoor air quality, classification of indoor air contaminations, health effects due to poor IAQ and the method to the remediation of poor IAQ. Chapter 3 provides in details the method of development used in this study including indoor fungal preparation, optimization of indoor fungal growth, the bio-coating resistance test and inhibition performance of *A. niger*. Then, chapter 4 is the result and discussion on the optimum growth of indoor fungus via response surface methodology and analysis on the growth percentages of *A. niger* treatment. Finally, chapter 5 is the conclusion for the overall objectives setup and further works to be recommended in the future.



PTTAA
PERPUSTAKAAN TUNKU TUNJUNG

REFERENCES

- Ababutain, I. M. (2013). Effect of some ecological factors on the growth of *Aspergillus niger* and *Cladosporium sphaerospermum*. *American Journal of Applied Sciences*, 10(2), 159–163.
- Adan, O. C. G. (1994). *On the fungal defacement of interior finishes*. Eindhoven University of Technology: PH.D. Thesis.
- Alghamdi, M. A., Shamy, M., Redal, M. A., Khoder, M., Awad, A. H., & Elserougy, S. (2014). Microorganisms associated particulate matter: a preliminary study. *The Science of the Total Environment*, 479–480, 109–16.
- Akzonobel (2017). What are materials made of. *Paint*. Retrieved from 10 February, 2018 from <http://resources.schoolscience.co.uk/ICI/14-16/paints/paintch1pg1.html>
- American Society for Testing and Material (2012). *Standard Test Method for Determining the Resistance of Paint Films and Related Coatings to Fungal Defacement Accelerated Four-Week Agar Plate Assay*. United States: ASTM D5560-00.
- American Society of Heating, Refrigerating and Air-Conditioning Engineers (2010). *Thermal Environmental Conditions for Human Occupancy*. Atlanta, GA:ASHRAE 62.
- American Society of Heating, Refrigerating and Air-Conditioning Engineers (2010). *Thermal Environmental Conditions for Human Occupancy*. Atlanta, GA:ASHRAE 55.
- Andersen, B., Frisvad, J. C., Søndergaard, I., Rasmussen, I. S., & Larsen, L. S. (2011). Associations between fungal species and water-damaged building materials. *Applied and Environmental Microbiology*, 77(12).

- Andersson, M. A., Nikulin, M., Kõljalg, U., Andersson, M. C., Rainey, F., & Reijula, K., Salkinoja-Salonen, M. (1997). Bacteria, molds, and toxins in water-damaged building materials. *Applied and Environmental Microbiology*, 63(2), 387–393.
- Andrews, S., & Pitt, J. I. (1987). Further studies on the water relations of xerophilic fungi, including some halophiles. *Microbiology*, 133(2), 233–238.
- Anwar, Z., Gulfraz, M., & Irshad, M. (2014). Agro-industrial lignocellulosic biomass a key to unlock the future bio-energy: A brief review. *Journal of Radiation Research and Applied Sciences*, 7(2), 163–173.
- Aoudou, Y., Leopold, T. N., Michel, J. D. P., Xavier, E. F., & Moses, M. C. (2010). Antifungal properties of essential oils and some constituents to reduce foodborne pathogen. *Journal of Yeast and Fungal Research*, 1(1), 001–008.
- Asan, A., Okten, S. S., & Sen, B. (2010). Airborne and soilborne microfungi in the vicinity hamitabat thermic power plant in Kırklareli City (Turkey), their seasonal distributions and relations with climatological factors. *Environmental Monitoring and Assessment*, 164, 221–231.
- Ayanbimpe, G.M., Danjuma, W.S., & Okolo, M.O. (2012). Relationship between fungal contamination of indoor air and health of some resident in Jos. *Intech Open Science*. 1-19.
- Ayliffe, G. A., Collins, B. J., Lowbury, E. J., Babb, J. R., & Lilly, H. A. (1967). Ward floors and other surfaces as reservoirs of hospital infection. *The Journal of Hygiene*, 65(4), 515–536.
- Bansod, S., & Rai, M. (2008). Antifungal activity of essential oils from Indian medicinal plants against human pathogenic *Aspergillus fumigatus* and *A. niger*. *World Journal of Medical Sciences*, 3(2), 81–88.
- Barzegar, H., Azizi, M. H., Barzegar, M., & Hamidi-esfahani, Z. (2014). Effect of potassium sorbate on antimicrobial and physical properties of starch – clay nanocomposite films. *Carbohydrate Polymers*, 110, 26–31.
- Baxter, C. G., Jones, A. M., Webb, K., & Denning, D. W. (2011). Homogenisation of cystic fibrosis sputum by sonication - An essential step for *Aspergillus* PCR. *Journal of Microbiological Methods*, 85(1), 75–81.
- Beezhold, D. H., Green, B. J., Blachere, F. M., Schmechel, D., Weissman, D. N., Velickoff, D., Wilson, N. W. (2008). Prevalence of allergic sensitization to indoor fungi in West Virginia. In *Allergy and Asthma Proceedings* 29(29–

34).

- Bekker, M., Erich, S. J. F., Hermanns, S. P. M., Maris, M. P. F. H. L. Van, Huinink, H. P., & Adan, O. C. G. (2015). Quantifying discoloration caused by the indoor fungus *Penicillium rubens* on building material at controlled humidity. *Building and Environment*, 90, 60–70.
- Bellotti, N., & Romagnoli, R. (2014). Assessment of Zinc Salicylate as Antifouling Product for Marine Coatings, 8–13.
- Bellotti, N., Salvatore, L., Deyá, C., Del Panno, M. T., del Amo, B., & Romagnoli, R. (2013). The application of bioactive compounds from the food industry to control mold growth in indoor waterborne coatings. *Colloids and Surfaces B: Biointerfaces*, 104, 140–144.
- Bertron, A., Escadeillas, G., & Duchesne, J. (2004). Cement pastes alteration by liquid manure organic acids: Chemical and mineralogical characterization. *Cement and Concrete Research*, 34(10), 1823–1835.
- Beukes, N., & Pletschke, B. I. (2006). Effect of sulfur-containing compounds on *Bacillus cellulosome*-associated “CMCase” and “Avicelase” activities. *FEMS Microbiology Letters*, 264(2), 226–231.
- Bluyssen, P. M., Cox, C., Seppänen, O., Oliveira Fernandes, E., Clausen, G., Müller, B., & Roulet, C. A. (2003). Why, when and how do HVAC-systems pollute the indoor environment and what to do about it? The European AIRLESS project. *Building and Environment*, 38(2), 209–225.
- Bonetta, S. S., Bonetta, S. S., Mosso, S., Sampò, S., Carraro, E., Samp, S., & Carraro, E. (2010). Assessment of microbiological indoor air quality in an Italian office building equipped with an HVAC system. *Environmental Monitoring and Assessment*, 161(1–4), 473–483.
- Boyacioglu, H., Haliki, A., Ates, M., Guvensen, A., & Abaci, O. (2007). The statistical investigation on airborne fungi and pollen grains of atmosphere in Izmir-Turkey. *Environmental Monitoring and Assessment*, 135(1–3), 327–334.
- Branz, L. (2007). *Designing Quality Learning Spaces: Ventilation & Indoor Air Quality*. Retrieved December 01, 2013, from Ministry of Education: <http://www.minedu.govt.nz/~media/MinEdu/Files/EducationSectors/PrimarySecondary/PropertyToolbox/ModernLearning/VentilationIndoorAirQualityGuide.pdf>

- Brian, G., Kimberly, H., Flanders, W.D., & George, K., (2002). Profiles of airborne fungi in buildings and outdoor environment in the United States. *Appl. Environ. Microbiol* 68(4), 1743-1753.
- Brijwani, K., Oberoi, H. S., & Vadlani, P. V. (2010). Production of a cellulolytic enzyme system in mixed-culture solid-state fermentation of soybean hulls supplemented with wheat bran. *Process Biochemistry*, 45(1), 120–128.
- Brischke, C., & Rapp, A. O. (2008). Dose-response relationships between wood moisture content, wood temperature and fungal decay determined for 23 European field test sites. *Wood Science and Technology*, 42(6), 507–518.
- Buchan, R. M., Rijal, P., Sandfort, D., & Keefe, T. (2002). Evaluation of airborne dust and endotoxin in corn storage and processing facilities in Colorado. *International Journal of Occupational Medicine and Environmental Health*, 15(1), 57–64.
- Bush, R. K. (2008). Indoor allergens, environmental avoidance, and allergic respiratory disease. *Allergy and Asthma Proceedings*, 29(6), 575–579.
- Carmo, E. S., Lima, E. D. O., De Souza, E. L., & De Sousa, F. B. (2008). Effect of *Cinnamomum zeylanicum* Blume essential oil on the growth and morphogenesis of some potentially pathogenic *Aspergillus* species. *Brazilian Journal of Microbiology*, 39(1), 91–97.
- Chapman, J. S. (2003). Biocide resistance mechanisms. *International Biodeterioration and Biodegradation*, 51(2), 133–138.
- Chipley, J.R., (2005). Sodium benzoate and benzoic acid. in; davidson pm, sofos jn. *Antimicrobial in foods*. 11-48.
- Cho, S. J., Park, J. H., Kreiss, K., & Cox-Ganser, J. M. (2011). Levels of microbial agents in floor dust during remediation of a water-damaged office building. *Indoor Air*, 21(5), 417–426.
- Cho, S. J., Park, J. H., Kreiss, K., & Cox-Ganser, J. M. (2011). Levels of microbial agents in floor dust during remediation of a water-damaged office building. *Indoor Air*, 21(5), 417–426.
- Choi, Y., Joe I, Lee, S., and Oh, K. (2009). An experimental study on the ignition and emissions characteristics of wall papers. *Mechanical Science and Technology*. 23, 2839-2847
- Clarke, J., Johnstone, C., Kelly, N., McLean, R., Anderson, J., Rowan, N., & Smith, J. (1999). A technique for the prediction of the conditions leading to mould

- growth in buildings. *Building and Environment*, 34(4), 515–521.
- Clausen, C. A., & Yang, V. (2007). Protecting wood from mould, decay, and termites with multi-component biocide systems. *International Biodeterioration & Biodegradation*, 59(1), 20–24.
- Clausen, C. A., & Yang, V. W. (2003). Mold inhibition on unseasoned southern pine. *The International Research Group on Wood Preservation*, 1–9.
- Contant, S., Júnior, G. C., Machado, P. F. M. P. B., & Lona, L. M. F. (2010). Evaluation of the effect of dry-film biocides on paint film preservation using neural networks. *Brazilian Journal of Chemical Engineering*, 27(4), 643–651.
- Cwalina, B. (2008). Biodeterioration of Concrete. *Journal of Architecture Civil Engineering Environment*, 133–140.
- D’Orazio, M., Palladini, M., Aquilanti, L., & Clementi, F. (2009). Experimental evaluation of the growth rate of mould on finishes for indoor housing environments: Effects of the 2002/91/EC directive. *Building and Environment*, 44(8), 1668–1674.
- Department of Occupational Safety and Health (2010). *Industry Code of Practice on Indoor Air Quality*. Malaysia : DOSH.
- Deabes, M. M., Abou El-Soud, N. H., & Abou El-Kassem, L. T. (2011). In vitro inhibition of growth and aflatoxin B1 production of aspergillus flavus strain (ATCC 16872) by various medicinal plant essential oils. *Macedonian Journal of Medical Sciences*, 4(4), 345–350.
- Dedesko, S., & Siegel, J. A. (2015). Moisture parameters and fungal communities associated with gypsum drywall in buildings. *Microbiome*, 3(1), 71.
- Di Giulio, M., Grande, R., Di Campli, E., Di Bartolomeo, S., & Cellini, L. (2010). Indoor air quality in university environments. *Environmental Monitoring and Assessment*, 170(1–4), 509–517.
- Dias, F., Mossini, S. A. G., M., Arrotéia, CC., da Costa, C. L., Nakamura, C. V., & Machinski, M. (2013). The inhibitory effects of Curcuma longa L. essential oil and curcumin on Aspergillus flavus link growth and morphology. *The Scientific World Journal*, 2013(2), 343804.
- Došen, I., Nielsen, K. F., Clausen, G., & Andersen, B. (2016). Potentially harmful secondary metabolites produced by indoor Chaetomium species on artificially and naturally contaminated building materials. *Indoor Air*, 1–13.
- Douwes, J., Thorne, P., Pearce, N., & Heederik, D. (2003). Bioaerosol health effects

- and exposure assessment: Progress and prospects. *Annals of Occupational Hygiene*, 47(3), 187–200.
- Dutkiewicz, J., Jabloski, L., & Olenchock, S. A. (1988). Occupational biohazards: A review. *American Journal of Industrial Medicine*, 14(5), 605–623.
- El-mougy, N. S., El-gamal, N. G., & Fotouh, Y. O. (2006). Evaluation of different application methods of chitin and chitosan for controlling tomato root rot disease under greenhouse and field conditions. *Egypt. J. Phytopathol*, 2(5), 190–195.
- Er, C. M., Sunar, N. M., Leman, A. M., Othman, N., Gani, P., Jamal, N. A., Parjo, U. K. (2015). Inhibitory Assay of an isolated indoor airborne fungus from an Institutional Building of Computer Education by using potassium sorbate. *Applied Mechanics and Materials*, 773–774, 1091–1095.
- Er, C. M., Sunar, N. M., Leman, A. M., & Othman, N. (2015). Direct growth inhibition assay of total airborne fungi with application of biocide-treated malt extract agar. *MethodsX*, 2, 340–344.
- Erich, S. J. F., Mendoza, S. M., Floor, W., Hermanns, S. P. M., Homan, W. J., & Adan, O. C. G. (2011). Decreased bio-inhibition of building materials due to transport of biocides. *Heron*, 56(3), 93–106.
- European Food Safety Authority (2012) Scientific opinion on the safety and efficacy of potassium sorbate for dogs and cats. Europe; EFSA 2012
- Fagundes, C., Pérez-Gago, M. B., Monteiro, A. R., & Palou, L. (2013). Antifungal activity of food additives in vitro and as ingredients of hydroxypropyl methylcellulose-lipid edible coatings against *Botrytis cinerea* and *Alternaria alternata* on cherry tomato fruit. *International Journal of Food Microbiology*, 166(3), 391–398.
- Favero, M. S., McDade, J. J., Robertsen, J. A., Hoffman, R. K., & Edwards, R. W. (1968). Microbiological Sampling of Surfaces. *Journal of Applied Bacteriology*, 31(3), 336–343.
- Fitzpatrick, C. S., Fradin, E., & Gregory, J. (2004). Temperature effects on flocculation, using different coagulants. *Water Science and Technology*, 50, 171–175.
- Flaherty, D. K., Deck, F. H., Hood, M.A., Liebert, C., Singleton, F., Winzenburger, P., Witmer, W. B. (1984). A *Cytophaga* species endotoxin as a putative agent of occupation-related lung disease. *Infection and Immunity*, 43(1), 213–216.

- Flannigan, B. & Miller, J.D (2011). Microbial growth in indoor environments. *Diversity, Health Impacts, Investigation and Control*. New York: CRC Press LLC.
- Flores-Cervantes, D. X., Palou, E., & Lopez-Malo, A. (2013). Efficacy of individual and combined UVC light and food antimicrobial treatments to inactivate *Aspergillus flavus* or *A. niger* spores in peach nectar. *Innovative Food Science and Emerging Technologies*, 20, 244–252.
- Frankel, M., Hansen, E. W., & Madsen, A. M. (2014). Effect of relative humidity on the aerosolization and total inflammatory potential of fungal particles from dust-inoculated gypsum boards. *Indoor Air*, 24(1), 16–28.
- Friman, J. (2010). *Comparative Study on Mould Growth on Plaster Boards treated with Biocides*. University of Goteborg: Degree Master Thesis.
- Gaskin, S., Taylor, M., Bentham, R., & Pisaniello, D. (2012). Understanding and managing risks associated with fungal contamination in indoor environments. University of Adelaide: Master's Thesis.
- Ghafoori, N., & Mathis, R. (1997). Sulfate resistance of concrete pavers. *Journal of Materials in Civil Engineering*, (February), 35–40.
- Giannantonio, D. J., Kurth, J. C., Kurtis, K. E., & Sobecky, P. a. (2009). Effects of concrete properties and nutrients on fungal colonization and fouling. *International Biodeterioration and Biodegradation*, 63(3), 252–259.
- Goly, C., Soro, Y., Kassi, B., Dadié, A., Soro, S., & Dje, M. (2015). Antifungal activities of the essential oil extracted from the tea of savanna (*Lippia multiflora*) in Côte d ' Ivoire, 9(February), 24–34.
- Gordon, T. (1992). Acute respiratory effects of endotoxin-contaminated machining fluid aerosols in Guinea pigs. *Toxicological Sciences*, 19(1), 117–123.
- Gots, R. E., Layton, N. J., & Pirages, S. W. (2010). Indoor health: background levels of fungi. *AIHA Journal : A Journal for the Science of Occupational and Environmental Health and Safety*, 64(4), 427–438.
- Gottschalk, C., Bauer, J., & Meyer, K. (2008). Detection of satratoxin G and H in indoor air from a water-damaged building. *Mycopathologia*, 166(2), 103–107.
- Grant, C., Hunter, C. A., Flannigan, B., & Bravery, A.F. (1989). The moisture requirements of moulds isolated from domestic dwellings. *International Biodeterioration*, 25(4), 259–284.
- Gregori, R., Borsetti, F., Neri, F., Mari, M., & Bertolini, P. (2008). Effects of

- potassium sorbate on postharvest brown rot of stone fruit. *Journal of Food Protection*, 71(8), 1626–1631.
- Guynot, M. E., Ramos, A. J., Sanchis, V., & Marín, S. (2005). Study of benzoate, propionate, and sorbate salts as mould spoilage inhibitors on intermediate moisture bakery products of low pH (4.5-5.5). *International Journal of Food Microbiology*, 101(1–2), 161–168.
- Khan, H., & Karuppayil, S. (2012). Fungal pollution of indoor environments and its management. *Saudi Journal of Biological Sciences*, 19(4), 405–426.
- Haliki-Uztan, A., Ateş, M., Abacı, Ö., Gülbahar, O., Erdem, N., Çiftçi, Ö., & Boyacıoğlu, H. (2010). Determination of potential allergenic fungal flora and its clinical reflection in suburban elementary schools in Izmir. *Environmental Monitoring and Assessment*, 168(1–4), 691–702.
- Halios, C. H., & Helmis, C. G. (2010). Temporal evolution of the main processes that control indoor pollution in an office microenvironment: A case study. *Environmental Monitoring and Assessment*, 167(1–4), 199–217.
- Halstensen, A. S. (2008). Species-specific fungal DNA in airborne dust as surrogate for occupational mycotoxin exposure. *International Journal of Molecular Sciences*, 9(12), 2543–2558.
- Hedayati, M. T., Mayahi, S., & Denning, D. W. (2010). A study on *Aspergillus* species in houses of asthmatic patients from Sari City, Iran and a brief review of the health effects of exposure to indoor *Aspergillus*. *Environmental Monitoring and Assessment*, 168(1–4), 481–487.
- Hervieux, V., Yaganza, E. S., Arul, J., & Tweddell, R. J. (2002). Effect of organic and inorganic salts on the development of *Helminthosporium solani*, the causal agent of potato silver scurf. *Plant Disease*, 86(9), 1014–1018.
- Heydaryinia, A., Veissi, M., & Sadadi, A. (2011). A comparative study of the effects of the two preservatives, sodium benzoate and potassium sorbate on *Aspergillus niger* and *Penicillium notatum*. *Jundishapur Journal of Microbiology*, 4(4), 301–307.
- Hoang, C. P., Kinney, K. a., Corsi, R. L., & Szaniszló, P. J. (2010). Resistance of green building materials to fungal growth. *International Biodeterioration & Biodegradation*, 64(2), 104–113.
- Hochmannova, L., & Vytrasova, J. (2010). Photocatalytic and antimicrobial effects of interior paints. *Progress in Organic Coatings*, 67(1), 1–5.

- Hood, J. R., Burton, D. M., Wilkinson, J. M., & Cavanagh, H. M. A. (2010). The effect of *Leptospermum petersonii* essential oil on *Candida albicans* and *Aspergillus fumigatus*. *Medical Mycology*, 48(7), 922–931.
- Hospodsky, D., Yamamoto, N., Nazaroff, W. W., Miller, D., Gorthala, S., & Peccia, J. (2015). Characterizing airborne fungal and bacterial concentrations and emission rates in six occupied children's classrooms. *Indoor Air*, 25(6), 641–652.
- Hyvärinen, A., Meklin, T., Vepsäläinen, A., & Nevalainen, A. (2002). Fungi and actinobacteria in moisture-damaged building materials - Concentrations and diversity. *International Biodeterioration and Biodegradation*, 49(1), 27–37.
- Ianovici, N., Maria, C., Răduțoiu, M. N., Haniș, A., & Tudorică, D. (2013). Variation in airborne fungal spore concentrations in four different microclimate regions in Romania. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 41(2), 450–457.
- Ibrahim, M., Rabah, A.B., Liman, B., & Ibrahim, N.T., (2011). Effect of temperature and relative humidity on the growth of *Helminthosporium fulvum*. *Nigerian Journal of Basic and Applied Science*. 19(1), 127-129.
- International Standard Organization (2005). *Ergonomics of the thermal environment: analytical determination and interpretation of thermal comfort using calculation of the PMV and PPD indices and local thermal comfort criteria*. Switzerland : ISO 7730.
- Iqbal, H. M. N. (2012). Enhanced bio-catalytic and tolerance properties of an indigenous cellulase through xerogel immobilization. *Advances in Bioscience and Biotechnology*, 3(4), 308–313.
- Iqbal, H. M. N., Ahmed, I., Zia, M. A., & Irfan, M. (2011). Purification and characterization of the kinetic parameters of cellulase produced from wheat straw by *Trichoderma viride* under SSF and its detergent compatibility. *Advances in Bioscience and Biotechnology*, 2(3), 149–156.
- Isaksson, T., Thelandersson, S., Ekstrand-Tobin, A., & Johansson, P. (2010). Critical conditions for onset of mould growth under varying climate conditions. *Building and Environment*, 45(7), 1712–1721.
- Ismail, S.H., Baba, D., & Abd Mutalib, L. (2010). Indoor air quality issues for non-industrial work place. *International Journal of Research and Reviews in Applied Sciences*, 5(3), 235–244.

- Siti, I., Baba, D., & Abd Mutalib, L. (2010). Indoor air quality issues for non-industrial work place. *International Journal of Research and Reviews in Applied Sciences*, 5(3), 235–244.
- Johansson, P., Tobin, A., Svensson, T., & Bok, G. (2012). Laboratory study to determine the critical moisture level for mould growth on building materials. *International Biodeterioration and Biodegradation*, 73, 23–32.
- Johansson, P., Samuelson, I., Ekstrand-tobin, A., Mjornell, K., Sandberg, P. I., & Sikander, E. (2005). *Microbiological growth on building materials – critical moisture levels . State of the art*.
- Johansson, P., Svensson, T., & Tobin, A. (2013). Validation of critical moisture conditions for mould growth on building materials. *Building and Environment*, 62, 201–209.
- Jones, A. P. (1999). Indoor air quality and health. *Atmospheric Environment*, 33(28), 4535–4564.
- Junqueira-Gonçalves, M. P., Alarcón, E., & Niranjana, K. (2016). The efficacy of potassium sorbate-coated packaging to control postharvest gray mold in raspberries, blackberries and blueberries. *Postharvest Biology and Technology*, 111, 205–208.
- Jussila, J., Komulainen, H., Huttunen, K., Roponen, M., Iivanainen, E., Torkko, P., Hirvonen, M.-R. (2002). Mycobacterium terrae isolated from indoor air of a moisture-damaged building induces sustained biphasic inflammatory response in mouse lungs. *Environmental Health Perspectives*, 110(11), 1119–25.
- Kampa, M., and Castanas, E., (2007). Human health effects of air pollution. *Environmental Pollution*.151, 362-367.
- Karunasena, E., Larraga, M. D., Simoni, J. S., Douglas, D. R., & Straus, D. C. (2010). Building-associated neurological damage modeled in human cells: a mechanism of neurotoxic effects by exposure to mycotoxins in the indoor environment. *Mycopathologia*, 170(6), 377–390.
- Kaufmann, S. H. E., & van Embden, J. D. A. (1993). Tuberculosis: a neglected disease strikes back. *Trends in Microbiology*, 1(1), 2–5.
- KEMI. (2012, March). *Biocide Treated Articles-An Internet Survey*. Retrieved November 13, 2013, from Swedish Chemicals Agency: www.kemi.se
- Khan, A. A. H., Karuppayil, S. M., Manoharachary, C., Kunwar, I. K., & Waghray,

- S. (2009). Isolation, identification and testing for allergenicity of fungi from air-conditioned indoor environments. *Aerobiologia*, 25(2), 119–123.
- Khan, M. S. A., & Ahmad, I. (2011). In vitro antifungal, anti-elastase and anti-keratinase activity of essential oils of Cinnamomum-, Syzygium- and Cymbopogon-species against *Aspergillus fumigatus* and *Trichophyton rubrum*. *Phytomedicine*, 19(1), 48–55.
- Khrueayu, D., and Pilantanapak, A., (2012). Antifungal activity of bioactive compound from endophytic fungi isolated from mangrove leaves. *Microbiology*. 1-6.
- Kinderlerer, J. L., & Hatton, P. V. (1990). Fungal metabolites of sorbic acid. *Food Additives and Contaminants*, 7(5), 657–669.
- Kishore, G. K., Pande, S., & Harish, S. (2007). Evaluation of essential oils and their components for broad-spectrum antifungal activity and control of late leaf spot and crown rot diseases in peanut. *Plant Disease*, 91(17), 375–379.
- Koc, F., & Kara, S. (2014). Environmental factors affecting efficacy of some essential oils and potassium sorbate to control growth of *Aspergillus flavus*, *Aspergillus parasiticus* on wheat and maize grains. *Journal of Agricultural Science and Technology*, 16(6), 1325–1334.
- Kontogeorgos, D. A., & Founti, M. A. (2013). A generalized methodology for the definition of reactive porous materials physical properties: Prediction of gypsum board properties. *Construction and Building Materials*, 48, 804–813.
- Kuo, N. W., Chiang, H. C., & Chiang, C. M. (2008). Development and application of an integrated indoor air quality audit to an international hotel building in Taiwan. *Environmental Monitoring and Assessment*, 147(1–3), 139–147.
- Lajili, H., Devillers, P., Grambin-Lapeyre, C., & Bournazel, J. P. (2008). Alteration of a cement matrix subjected to biolixiviation test. *Materials and Structures*, 41(10), 1633–1645.
- Lander, F., Meyer, H. W., & Norn, S. (2001). Serum IgE specific to indoor moulds, measured by basophil histamine release, is associated with building-related symptoms in damp buildings. *Inflammation Research*, 50(4), 227–231.
- Lanier, C., Richard, E., Heutte, N., Picquet, R., Bouchart, V., & Garon, D. (2010). Airborne molds and mycotoxins associated with handling of corn silage and oilseed cakes in agricultural environment. *Atmospheric Environment*, 44(16), 1980–1986.
- Laverge, J., Van Den Bossche, N., Heijmans, N., & Janssens, A. (2011). Energy

- saving potential and repercussions on indoor air quality of demand controlled residential ventilation strategies. *Building and Environment*, 46(7), 1497–1503.
- Leona, K., and Hawks (1995). *Home furnishings and housing specialist*. Wood Finishing & Refinishing, 2(3), 173-80.
- Lewinska, A. M., Hoof, J. B., Peuhkuri, R. H., Rode, C., Lilje, O., Foley, M., ... Andersen, B. (2016). Visualization of the structural changes in plywood and gypsum board during the growth of *Chaetomium globosum* and *Stachybotrys chartarum*. *Journal of Microbiological Methods*, 129, 28–38.
- Li, X. F., & Feng, X. Q. (2008). Effects of molecularweight and concentration of chitosan on antifungal activity against *Aspergillus Niger*. *Iranian Polymer Journal*, 17(Rew98), 843–852.
- Lian, K. S., Sri, P., Inangda, N., Ramly, A., & Inangda, P. S. N. (2007). Sources of Indoor Air Quality Problem in a New Hospital in Malaysia. *The 6th International Conference on Indoor Air Quality, Ventilation & Energy Conservation in Buildings, Sendai, Japan*.
- Loftness, V., Hakkinen, B., Adan, O., & Nevalainen, A. (2007). Elements that contribute to healthy building design. *Environmental Health Perspectives*, 115(6), 965–970.
- López, O. V., Giannuzzi, L., Zaritzky, N. E., & García, M. A. (2013). Potassium sorbate controlled release from corn starch films. *Materials Science and Engineering C*, 33(3), 1583–1591.
- Louthrenoo, E., Park, Y.S., Philippe, L., and Schumacher, H.R.(1990). Localized peripheral calcium oxalate crystal deposition caused by *Aspergillus Niger* infection. *Journal Rheumatol*, (17)407412.
- Luksamijarulkul, P., Ratthanakhot, Y. & Vatanasomboon, P. (2012). Microbial counts and particulate matter levels in indoor air samples collected from a Child Home-Care Center in Bangkok, Thailand. *J Med Assoc Thai* , 161-168.
- Madan, R. K., & Levitt, J. (2014). A review of toxicity from topical salicylic acid preparations. *Journal of the American Academy of Dermatology*, 70(4), 788–792.
- Mainelis, G., Grinshpun, S.A., Adhirakari, A. (2002).Design and collection efficiency of a new electrostatic precipitator for bioaerosol collection. *Aerosol Sci Technol*.36, 1073-85.

- Mandal, J., And Brandl, H., (2011). Bioaerosol in indoor environment – A review with special reference to residential and occupational locations. *Environmental & Biological Monitoring*, 4, 83-96
- Maroni, M., Seifert, B., Lindvall, T., (1995). *Indoor Air Quality*. Air Quality Monographs (3), 1049.
- Miller J.D., Rand T.G., Jarvis B.B., (2003). Stachybotrys. Medical: cause of human disease or media. *Mycology*, (4) 271-291
- Mehrara, E., Forssell-Aronsson, E., Ahlman, H., & Bernhardt, P. (2009). Quantitative analysis of tumor growth rate and changes in tumor marker level: specific growth rate versus doubling time. *Acta Oncologica (Stockholm, Sweden)*, 48(4), 591–597.
- Mohd, N., Noraini, R., Leman, A. M., Sayuti, A., & Abidin, Z. (2013). Building and indoor environment : a study on three stages of a new building commissioning, 3(3), 63–68.
- Murtoniemi, T., Nevalainen, A., & Hirvonen, M. R. (2003). Effect of plasterboard composition on Stachybotrys chartarum growth and biological activity of spores. *Applied and Environmental Microbiology*, 69(7), 3751–3757.
- Nalisha I., Muskhazli M., and Farizan T., (2006). *Production of bioactive compounds by bacillus subtilis against sclerotium rolfsii*. 2(2), 19-23.
- National Institute of Occupational Safety & Health (2010). *Recommendations for the Cleaning and Remediation of Flood-Contaminated HVAC Systems: A Guide for Building Owners and Managers*. Retrieved from 9 June, 2013 from: <http://www.cdc.gov/niosh/topics/emres/Cleaning-Flood-HVAC.html>.
- Nasir, Z. A., Colbeck, I., Sultan, S., & Ahmed, S. (2012). Bioaerosols in residential micro-environments in low income countries: A case study from Pakistan. *Environmental Pollution*, 168, 15–22.
- Nawar, L. S. (2008). Prevention and control of fungi contaminated stored pistachio nuts imported to Saudi Arabia. *Journal of Biological Sciences*, 15(1), 105–112.
- Nazar, F. N., Weerasinghe, T., & Perera, S. (2013). A preliminary study on the resistance of paints against the growth of algae and fungi. *Advanced Materials Research*, 746, 186–191.
- Nevalainen, A., Täubel, M., & Hyvärinen, A. (2015). Indoor fungi: Companions and contaminants. *Indoor Air*, 25(2), 125–156.
- Nielsen, K. F. (2002). *Mould growth on building materials. Secondary metabolites*,

mycotoxins and biomarkers. Building.

- Nielsen, K. F., Holm, G., Uttrup, L. P., & Nielsen, P. A. (2004). Mould growth on building materials under low water activities. Influence of humidity and temperature on fungal growth and secondary metabolism. *International Biodeterioration and Biodegradation*, 54(4), 325–336.
- Niemenmaa, O. (2008). *Monitoring of fungal growth and degradation of wood*. Retrieved from <http://doria17-kk.lib.helsinki.fi/handle/10024/37369>.
- Nikolov, A., and Ganchev, D.,(2011). In vitro antifungal examination of potassium sorbate towards some phytopathogens. *Agricultural Academy*,17(2), 191-194.
- Nikulin, M., Pasanen, A. L., Berg, S., & Hintikka, E. L. (1994). Stachybotrys atra growth and toxin production in some building materials and fodder under different relative humidities. *Applied and Environmental Microbiology*, 60(9), 3421–3424.
- Norhidayah, A., Chia-Kuang, L., Azhar, M. K., & Nurulwahida, S. (2013). Indoor air quality and sick building syndrome in three selected buildings. *Procedia Engineering*, 53(2010), 93–98.
- Norback, D., Hashim, J. H., Hashim, Z., Cai, G. H., Sooria, V., Ismail, S. A., & Wieslander, G. (2017). Respiratory symptoms and fractional exhaled nitric oxide (FeNO) among students in Penang, Malaysia in relation to signs of dampness at school and fungal DNA in school dust. *Science of the Total Environment*, 577, 148–154.
- Norback, D., Hashim, J. H., Markowicz, P., Cai, G. H., Hashim, Z., Ali, F., & Larsson, L. (2016). Endotoxin, ergosterol, muramic acid and fungal DNA in dust from schools in Johor Bahru, Malaysia - Associations with rhinitis and sick building syndrome (SBS) in junior high school students. *Science of the Total Environment*, 545–546, 95–103.
- Nourah, A. P. (2008). Indoor Fungal and Bacterial Contaminations on Household Environment in Riyadh , Saudi Arabia(June), 15(1)113-119.
- Occupational Safety and Health Administration (2011). *Indoor Air Quality in Commercial and Institutional Buildings*. United States: OSHA 3430.
- OHS. (2012). *Indoor Air Quality (IAQ)*. Retrieved October 06, 2013, from Occupational Health and Safety Bulletin, Alberta: www.employment.alberta.ca/ohs-legislation.
- Ojanen, T., Viitanen, H., Peuhkuri, R., Vinha, J., & Salminen, K. (2010). Mold growth

modeling of building structures to improve the model.

- Palacios, H., Taniwaki, M. H., Hashimoto, J. M., & De Menezes, H. C. (2005). Growth of *Aspergillus ochraceus*, *A. carbonarius* and *A. niger* on culture media at different water activities and temperatures. *Brazilian Journal of Microbiology*, 36(1), 24–28.
- Panackal, A.A., Imhof, A., Hanley, E.W., Marr, K.A., (2006). *Aspergillus ustus* infections among transplant recipients. *Emerg Infect Dis.* 12, 403–408.
- Park, J. H., Cox-Ganser, J. M., White, S. K., Laney, A. S., Caulfield, S. M., Turner, W. A., Kreiss, K. (2016). Bacteria in a water-damaged building: Associations of actinomycetes and non-tuberculous mycobacteria with respiratory health in occupants. *Indoor Air*, 1–10.
- Pasanen, A.L., Kasanen, J.-P., Rautiala, S., Ikäheimo, M., Rantamäki, J., Kääriäinen, H., & Kalliokoski, P. (2000). Fungal growth and survival in building materials under fluctuating moisture and temperature conditions. *International Biodeterioration & Biodegradation*, 46(2), 117–127.
- Patrick, C. N., & Ebere, M. (2014). Optimization of chemical treatment conditions for *adenia lobata* fiber using CCD, 4(6), 23–32.
- Pawar, V. C., & Thaker, V. S. (2006). In vitro efficacy of 75 essential oils against *Aspergillus niger*. *Mycoses*, 49(4), 316–323.
- Penny, M., Etherton, K., Kari, D., Bonanome, A., (2002). Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Excerpta Medica, Inc.* 113(9B), 71–87.
- Pereira, A., Palha, F., Brito, J. D., & Silvestre, J. D. (2011). Inspection and diagnosis system for gypsum plasters in partition walls and ceilings. *Construction and Building Materials*, 25(4), 2146–2156.
- Pillai, P., & Ramaswamy, K. (2012). Effect of naturally occurring antimicrobials and chemical preservatives on the growth of *Aspergillus parasiticus*. *Journal of Food Science and Technology*, 49(2), 228–233.
- Pitkäranta, M., Meklin, T., Hyvärinen, A., Nevalainen, A., Paulin, L., Auvinen, P., Rintala, H. (2011). Molecular profiling of fungal communities in moisture damaged buildings before and after remediation a comparison of culture-dependent and culture-independent methods. *BMC Microbiology*, 11, 235.
- Portnoy, J. M. (2003). Evaluation of indoor mold exposure is what allergists do best. *Annals of Allergy, Asthma & Immunology: Official Publication of the*

American College of Allergy, Asthma, & Immunology, 90(2), 175.

- Rahman, W. A., Rosli, H., Baharuddin, S. N., & Salleh, B. (2012). Incidence and remediation of fungi in a sick building in Malaysia: A case study. *Aerobiologia*, 28(2), 275–283.
- Rakotonirainy, M. S., Vilmont, L. B., & Lavédrine, B. (2016). A methodology for detecting the level of fungal contamination in the French Film Archives vaults. *Journal of Cultural Heritage*, 19, 454–462.
- Rao, C. Y., Burge, H. a, & Chang, J. C. (1996). Review of quantitative standards and guidelines for fungi in indoor air. *Journal of the Air & Waste Management Association* (1995), 46(9), 899–908.
- Raskin, I. (1992). Role of salicylic acid in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 43(1), 439–463.
- Rasooli, I., Rezaei, M. B., & Allameh, A. (2006). Growth inhibition and morphological alterations of *Aspergillus niger* by essential oils from *Thymus eriocalyx* and *Thymus x-porlock*. *Food Control*, 17(5), 359–364.
- Ravikumar, H., Shwetha, S. R., and Karigar, C.S., (2012). *Biodegradation of Paints*, 5(1), 1977-1983.
- Reddy, M. K., & Srinivas, T. (2017). Mold Allergens in Indoor Play School Environment. *Energy Procedia*, 109 (November 2016), 27–33.
- Reeslev, M., Miller, M., & Nielsen, K. F. (2003). Quantifying Mold Biomass on Gypsum Board: Comparison of Ergosterol and Beta- N - Acetylhexosaminidase as Mold Biomass Parameters Quantifying Mold Biomass on Gypsum Board: Comparison of Ergosterol and Beta- N - Acetylhexosaminidase as Mold Biomass Paramete, 69(7), 5–8.
- Rode, C., Ph, D., Grau, K., & Mitamura, T. (n.d.). Model and experiments for hygrothermal conditions of the envelope and indoor air of buildings.
- Roden, K. (2010). Biocides in Antimicrobial Paints. *Microbiology Australia*, 198-200.
- Rutala, W. A., & Weber, D. J. (2004). Disinfection and sterilization in health care facilities: what clinicians need to know. *Clinical Infectious Diseases*, 39, 702–709.
- Ryu, S. H., & Moon, H. J. (2014). Mould germination and the growth rate on wallpapers with different physical properties and the surface structures. *Indoor and Built Environment*, 23(1), 171–179.
- Saari, S., Mensah-Attipoe, J., Reponen, T., Veijalainen, A. M., Salmela, A., Pasanen,

- P., & Keskinen, J. (2015). Effects of fungal species, cultivation time, growth substrate, and air exposure velocity on the fluorescence properties of airborne fungal spores. *Indoor Air*, 25(6), 653–661.
- Saleh M. Al-Garni, Saleh Kabli, F. A.-S. and Z. A.-G. (2007). Mycoflora Associated with Some Textiles in Jeddah Cuty, *Vol/19*(March), 99:93-113.
- Salem, M. Z. M., Zidan, Y. E., Mansour, M. M. a, El Hadidi, N. M. N., & Abo Elgat, W. A., A. (2016). Antifungal activities of two essential oils used in the treatment of three commercial woods deteriorated by five common mold fungi. *International Biodeterioration and Biodegradation*, 106, 88–96.
- Samson, R. A., Visagie, C. M., Houbaken, J., Hong, S. B., Hubka, V., Klaassen, C. H. W., Frisvad, J. C. (2014). Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology*, 78, 141–173.
- Santilli, J., & Rockwell, W. (2003). Fungal contamination of elementary schools: a new environmental hazard. *Annals of Allergy, Asthma & Immunology*. 90(2), 203–8.
- Sarbu, I., & Sebarchievici, C. (2013). Aspects of indoor environmental quality assessment in buildings. *Energy and Buildings*, 60, 410–419.
- Savkovi, Eljko D., Stupar, M., Grbi, M. V. L., & Vukojevi, J. B. (2016). Comparison of anti-*Aspergillus* activity of *Origanum vulgare* L. essential oil and commercial biocide based on silver ions and hydrogen peroxide. *Acta Botanica Croatica*, 75(1), 121–128.
- Schleibinger, H., Keller, R., & Roden, H. (2004). *Indoor Air Pollution by Microorganisms and Their Metabolites. Handbook of Environmental Chemistry* (Vol. 4).
- Schuster, E., Dunn-Coleman, N., Frisvad, J., & Van Dijck, P. (2002). On the safety of *Aspergillus niger* - A review. *Applied Microbiology and Biotechnology*, 59(4–5), 426–435.
- Sen, B., & Asan, A. (2009). Fungal flora in indoor and outdoor air of different residential houses in Tekirdag City (Turkey): Seasonal distribution and relationship with climatic factors. *Environmental Monitoring and Assessment*, 151(1–4), 209–219.
- Shahmohammadi, M., Javadi, M., & Nassiri-Asl, M. (2016). An Overview on the effects of Sodium Benzoate as a preservative in food products. *Biotechnology and Health Sciences*, 3(3).

- Shelton, B. G., Kirkland, K. H., Flanders, W. D., & Morris, G. K. (2002). Profiles of Airborne fungi in buildings and outdoor environments in the United States. *Applied and Environmental Microbiology*, 68(4), 1743–1753.
- Shen, H., Tan, H., & Tzempelikos, A. (2011). The effect of reflective coatings on building surface temperatures, indoor environment and energy consumption - An experimental study. *Energy and Buildings*, 43(2–3), 573–580.
- Shirakawa, M. a., Gaylarde, C. C., Sahão, H. D., & Lima, J. R. B. (2013). Inhibition of Cladosporium growth on gypsum panels treated withnanosilver particles. *International Biodeterioration and Biodegradation*, 85, 57–61.
- Shirakawa, M. A., Gaylarde, C. C., Gaylarde, P. M., John, V., & Gambale, W. (2002). Fungal colonization and succession on newly painted buildings and the effect of biocide. *FEMS Microbiology Ecology*, 39(2), 165–173.
- Shirakawa, M. A., Selmo, S. M., Cincotto, M. a., Gaylarde, C., Brazolin, S., & Gambale, W. (2002). Susceptibility of phosphogypsum to fungal growth and the effect of various biocides. *International Biodeterioration and Biodegradation*, 49(4), 293–298.
- Singh, J., Yu, C., & Jeong Tai Kim. (2011). Building pathology toxic mould remediation. *Indoor and Built Environment*, 20(1), 36–46.
- Singh, T., & Chittenden, C. (2010). Efficacy of essential oil extracts in inhibiting mould growth on panel products. *Building and Environment*, 45(10), 2336–2342.
- Siripong, S., & Rittmann, B. E. (2007). Diversity study of nitrifying bacteria in full-scale municipal wastewater treatment plants. *Water Research*, 41(5), 1110–1120.
- Sivasubramani, S. K., Niemeier, R. T., Reponen, T., & Grinshpun, S. A. (2004). Assessment of the aerosolization potential for fungal spores in moldy homes. *Indoor Air*, 14(6), 405–412.
- Srinivasan, M., Natarajan, K., & Nagarajan, G. (2000). Growth optimization of an ectomycorrhizal fungus with respect to pH and temperature in vitro, using Design of Experiments. *Bioprocess Engineering*, 22(3), 0267–0273.
- Stanojevic, D., Comic, L., Stefanovic, O., & Solujic-Sukdolac, S. (2009). Antimicrobial effects of sodium benzoate, sodium nitrite and potassium sorbate and their synergistic action in vitro. *Bulgarian Journal of Agricultural Science*, 15(4), 307–311.

- Sundell, J. (2004). On the history of indoor air quality and health. *Indoor Air*, 14 Suppl 7(Suppl 7), 51–58.
- Sundis, M.S.A., Baharuddin, S., Sulaiman, A.K. & Shaida, F.S. (2012). Antimicrobial activity of spermine alkaloids from samanea saman against microbes associated with sick buildings. *International Conference on Environment, Chemistry and Biology*, 150-155.
- Syazwan, A. I., Juliana, J., Norhafizalina, O., Azman, Z. A, & Kamaruzaman, J. (2009). Indoor air quality and sick building syndrome in malaysian buildings. *Global Journal of Health Science*, 1(2), 126–135.
- Thermal Environmental Conditions for Human Occupancy*. Atlanta, GA: ASHRAE Standard 55-1992.
- Todero Ritter, C. E., Camassola, M., Zampieri, D., Silveira, M. M., & Dillon, A. J. P. (2013). Cellulase and xylanase production by penicillium echinulatum in submerged media containing cellulose amended with sorbitol. *Enzyme Research*, 2013, 1-9.
- Tortorano, A. M., Viviani, M. A., Biraghi, E., Rigoni, A. L., Prigitano, A., & Grillot, R. (2005). In vitro testing of fungicidal activity of biocides against *Aspergillus fumigatus*. *Journal of Medical Microbiology*, 54(10), 955–957.
- Tripti, S., and Chittenden, C., (2010). Efficacy of essential oil extract in inhibiting mould growth on panel products. *Building and environment*. (45)2336-2342.
- TSI Incorporated (2011). *A practical guide to indoor air quality investigations*. United States, Indoor air quality handbook.
- Türe, H., Gällstedt, M., & Hedenqvist, M. S. (2012). Antimicrobial compression-moulded wheat gluten films containing potassium sorbate. *Food Research International*, 45(1), 109–115.
- United States EPA (2010 April). *Indoor Air Pollution: An Introduction for Health Professionals*. Retrieved on 18 April 2013, from <http://www.epa.gov/iaq/pubs/hpguide.html>
- United States Environmental Protection Agency (2013). *A brief Guide to Mold, Moisture and your Home*. United States: EPA
- Vacher, S., Hernandez, C., Bärtschi, C., & Poussereau, N. (2010). Impact of paint and wall-paper on mould growth on plasterboards and aluminum. *Building and Environment*, 45(4), 916–921.
- Valerio, F., Favilla, M., De Bellis, P., Sisto, A., de Candia, S., & Lavermicocca, P.

- (2009). Antifungal activity of strains of lactic acid bacteria isolated from a semolina ecosystem against *Penicillium roqueforti*, *Aspergillus niger* and *Endomyces fibuliger* contaminating bakery products. *Systematic and Applied Microbiology*, 32(6), 438–448.
- Van Hoek, P., Van Dijken, J. P., & Pronk, J. T. (1998). Effect of specific growth rate on fermentative capacity of baker's yeast. *Applied and Environmental Microbiology*, 64(11), 4226–33.
- Varani, S., Stanzani, M., Paolucci, M., Melchionda, F., Castellani, G., Nardi, L., Sambri, V. (2009). Diagnosis of bloodstream infections in immunocompromised patients by real-time PCR. *Journal of Infection*, 58(5), 346–351.
- Verdier, T., Coutand, M., Bertron, A., & Roques, C. (2014). A review of indoor microbial growth across building materials and sampling and analysis methods. *Building and Environment*, 80, 136–149.
- Viitanen, H. (1994). Factors affecting the development of biodeterioration in wooden constructions. *Materials and Structures*, 27(8), 483–493.
- Viitanen, H., & Salonvaara, M. (1806). Moisture conditions and biodeterioration risk of building materials and structure. *Indoor Air*, (Figure 1), 1001–1006.
- Vojdani, A., Kashanian, A., Vojdani, E., & Campbell, A. W. (2003). Saliva secretory IgA antibodies against molds and mycotoxins in patients exposed to toxigenic fungi. *Immunopharmacology and Immunotoxicology*, 25(4), 595–614.
- Wackerl, A., Groß, A., & Hurraß, J. (2013). Isolation of fungal DNA from building materials. *Umweltmedizin in Forschung Und Praxis*, 18(5), 275–283.
- Walker, C. M., Abbott, G. F., Greene, R. E., Shepard, J. A. O., Vummidi, D., & Digumarthy, S. R. (2014). Imaging pulmonary infection: Classic signs and patterns. *American Journal of Roentgenology*, 202(3), 479–492.
- Wang, C. C., Fang, G. C., & Kuo, C. H. (2010). Bioaerosols as contributors to poor air quality in Taichung City, Taiwan. *Environmental Monitoring and Assessment*, 166(1–4), 1–9.
- Wang, J., & Oh, D. H. (2012). Effect of temperature and relative humidity on growth behavior of *Escherichia Coli* O157:H7 on Spinach using response surface methodology. *Journal of Food Safety*, 32(3), 296–304.
- Warscheid, T., & Braams, J. (2000). Biodeterioration of stone: A review. *International Biodeterioration and Biodegradation*, 46(4), 343–368.

- Wessen, B. (2006). The microbial flora in moisture damaged building material from swedish problem buildings. *Healthy Buildings*, (2006), 407-410.
- Wilson, S. C., Carriker, C. G., Brasel, T. L., Karunasena, E., Douglas, D. R., Wu, C., Straus, D. C. (2004). Culturability and toxicity of sick building syndrome-related fungi over time. *Journal of Occupational and Environmental Hygiene*, 1(8), 500–504.
- World Health Organization (2009). *WHO Guidelines for Indoor Air Quality: Dampness and Mould*. Retrieved from 26 May, 2013 from http://www.euro.who.int/__data/assets/pdf_file/0017/43325/E92645.
- Xiao F.L., Xiao Q.F., and Sheng Y. (2008). Effects of molecular weight and concentration of chitosan on antifungal activity against aspergillus niger. *Iranian Polymer Journal*, 17, 843–852.
- Yamamoto, N., Hospodsky, D., Dannemiller, K. C., Nazaroff, W. W., & Peccia, J. (2015). Indoor emissions as a primary source of airborne allergenic fungal particles in classrooms. *Environmental Science and Technology*, 49(8), 5098–5106.
- Yassin, M. F., and Atmougatea, S., (2010). *Assessment of airborne bacteria and fungi in an outdoor and outdoor environment*, 7(3), 535-544.
- York, G. K., & Vaughn, R. H. (1964). Mechanisms in the Inhibition of Microorganisms By Sorbic Acid. *Journal of Bacteriology*, 88(2), 411–417.
- Yusof, M. Y. M., Deros, B. M., & Ismail, A. R. (2013). Effect of temperature, humidity and illuminance towards worker's performance in automotive industry. *Sains Malaysiana*, 42(12), 1815–1818.
- Zabka, M., Pavela, R., & Prokinova, E. (2014). Antifungal activity and chemical composition of twenty essential oils against significant indoor and outdoor toxigenic and aeroallergenic fungi. *Chemosphere*, 112, 443–448.
- Zielecka, M., Bujnowska, E., Kępska, B., Wenda, M., & Piotrowska, M. (2011). Antimicrobial additives for architectural paints and impregnates. *Progress in Organic Coatings*, 72(1–2), 193–201.
- Zucker, B. A., Trojan, S., & Müller, W. (2000). Airborne gram-negative bacterial flora in animal houses. *Journal of Veterinary Medicine, Series B*, 47(1), 37–46.